

Complexity of the Two-Variable Fragment with (Binary Coded) Counting Quantifiers

Ian Pratt-Hartmann
Department of Computer Science,
University of Manchester.

Abstract

We show that the satisfiability and finite satisfiability problems for the two-variable fragment of first-order logic with counting quantifiers are both in NEXPTIME, even when counting quantifiers are coded succinctly.

1 Background

The two-variable fragment with counting quantifiers, here denoted $\mathcal{C}^2$, is the set of function-free, first-order formulas containing at most two variables, but with the counting quantifiers $\exists_{\leq C}$, $\exists_{\geq C}$ and $\exists_{=C}$ (for every $C > 0$) allowed. The *satisfiability problem*, $\text{Sat-}\mathcal{C}^2$, is the problem of deciding, for a given sentence ϕ of $\mathcal{C}^2$, whether ϕ has a model; the *finite satisfiability problem*, $\text{Fin-Sat-}\mathcal{C}^2$, is the problem of deciding, for a given sentence ϕ of $\mathcal{C}^2$, whether ϕ has a finite model. It is well-known that $\mathcal{C}^2$ lacks the finite model property; hence $\text{Sat-}\mathcal{C}^2$ and $\text{Fin-Sat-}\mathcal{C}^2$ do not coincide. The decidability of $\text{Sat-}\mathcal{C}^2$ was shown independently by Grädel, Otto and Rosen [2], and by Szwaart and Tendera [4, 5]. For a more general survey and some recent extensions, see Grädel and Otto [1] and Otto [3].

When discussing the computational complexity of these problems, it is important to specify how the sizes of numerical quantifier subscripts are measured. Under *unary coding*, a counting subscript C is taken to have size C ; under *binary coding*, by contrast, the same subscript is taken to have size $\log C$. In determining upper complexity-bounds, unary coding is evidently the less stringent accounting method, because the size of formulas involving counting quantifiers is, in general, exponentially larger than under binary coding. Pacholski *et al.*, *op. cit.* showed that $\text{Sat-}\mathcal{C}^2$ is in NEXPTIME, but only under the assumption of unary coding. The present paper shows that both $\text{Fin-Sat-}\mathcal{C}^2$ and $\text{Sat-}\mathcal{C}^2$ are in NEXPTIME, even with binary coding. We remark that both $\text{Fin-Sat-}\mathcal{C}^2$ and $\text{Sat-}\mathcal{C}^2$ are easily shown to be NEXPTIME-hard on either coding scheme.

In the sequel, we confine attention to finite or countably infinite structures interpreting finite signatures of unary and binary predicates, without individual constants or function-symbols; in addition, we treat the equality predicate $\approx$ as a

logical constant. The lack of individual constants in the signature constitutes no essential restriction of expressive power, because their effect can be reproduced using formulas of the form $\exists_{=1}xp(x)$. The presence of equality in the logic constitutes no extension of expressive power, because it can be defined by the formula $\forall x(x \approx x) \wedge \forall x\exists_{=1}y(x \approx y)$.

2 Preliminaries

If ϕ is a formula, let $\|\phi\|$ denote the number of symbols in ϕ , assuming that a numerical quantifier subscript C involves $\log C$ symbols. Thus, if the quantifier $\exists_{=C}$ occurs in ϕ , we have $C \leq 2^{\|\phi\|}$. The following type of normal-form lemma is standard in treatments of $\mathcal{C}^2$. (Recall that, in this paper, we are confining attention to signatures with no individual constants or function-symbols.)

Lemma 1. *Let ϕ be a sentence in $\mathcal{C}^2$. We can construct, in time bounded by a polynomial function of $\|\phi\|$, a formula*

$$\phi^* := \forall x\alpha \wedge \forall x\forall y(\beta \vee x \approx y) \wedge \bigwedge_{1 \leq h \leq m} \forall x\exists_{=C_h}y(f_h(x, y) \wedge x \not\approx y)$$

satisfying the following conditions: (i) α is a quantifier-free, equality-free formula with x as its only variable, (ii) β is a quantifier-free, equality-free formula with x and y as its only variables, (iii) the f_h ($1 \leq h \leq m$) are binary predicates, (iv) ϕ^* is satisfiable if and only if ϕ is satisfiable over a domain of size greater than $C = \max_{1 \leq h \leq m} C_h$.

Proof. Routine adaptation of textbook transformation to Scott normal form. See, for example, Grädel and Otto [1] Sec. 2.1 for an explanation of the required techniques. $\square$

We remark that, since C is bounded by a singly exponential function of $\|\phi\|$, satisfiability of ϕ over domains of size C or less can certainly be determined in NEXPTIME.

We next review some standard concepts. Fix a finite signature Σ of unary and binary predicates (no individual constants or function-symbols). A *literal* is an atomic formula or the negation of an atomic formula. A *1-type* is a maximal consistent set of (equality-free) literals involving only the variable x . A *2-type* is a maximal consistent set of equality-free literals involving only the variables x and y . If τ is a 2-type, then the result of transposing the variables x and y in τ will also be a 2-type, denoted τ^{-1} . If $\mathfrak{A}$ is any structure interpreting Σ , and $a \in A$, then there exists a unique 1-type $\pi(x)$ such that $\mathfrak{A} \models \pi[a]$; we denote π by $\text{tp}^{\mathfrak{A}}[a]$. If, in addition, $b \in A$ is distinct from a , then there exists a unique 2-type $\tau(x, y)$ such that $\mathfrak{A} \models \tau[a, b]$; we denote τ by $\text{tp}^{\mathfrak{A}}[a, b]$. We do not define $\text{tp}^{\mathfrak{A}}[a, b]$ if $a = b$. If π is a 1-type, we say that π is *realized* in $\mathfrak{A}$ if there exists $a \in A$ with $\text{tp}^{\mathfrak{A}}[a] = \pi$. If τ is a 2-type, we say that τ is *realized* in $\mathfrak{A}$ if there exist distinct $a, b \in A$ with $\text{tp}^{\mathfrak{A}}[a, b] = \tau$.

Notation 1. Any 2-type τ includes a unique 1-type, denoted $\text{tp}_1(\tau)$; in addition we write $\text{tp}_2(\tau)$ for $\text{tp}_1(\tau^{-1})$.

Remark 1. If $\text{tp}^{\mathfrak{A}}[a, b] = \tau$, then $\text{tp}^{\mathfrak{A}}[b, a] = \tau^{-1}$, $\text{tp}^{\mathfrak{A}}[a] = \text{tp}_1(\tau)$ and $\text{tp}^{\mathfrak{A}}[b] = \text{tp}_2(\tau)$.

We next introduce some non-standard concepts.

Definition 1. A *counting signature* Σ is a signature of unary and binary predicates (no individual constants or function-symbols) with a distinguished subset of binary predicates; we refer to these distinguished binary predicates as the *counting predicates* of Σ .

Counting signatures help us analyse models of formulas in $\mathcal{C}^2$ by allowing us to treat a subset of the binary predicates in a special way. Since counting signatures are signatures, we may meaningfully speak of 1-types and 2-types over counting signatures.

Definition 2. Let Σ be a counting signature, and let τ be a 2-type over Σ . We say that τ is a *message-type* (over Σ) if, for some counting predicate f of Σ , $f(x, y) \in \tau$. If τ is a message-type such that τ^{-1} is also a message-type, we say that τ is *invertible*. On the other hand, if τ is a 2-type such that neither τ nor τ^{-1} is a message-type, we say that τ is *silent*.

Thus, a 2-type τ is an invertible message-type if and only if there are counting predicates f and g such that $f(x, y) \in \tau$ and $g(y, x) \in \tau$. The terminology is meant to suggest the following imagery. If $\text{tp}^{\mathfrak{A}}[a, b]$ is a message-type μ , then we may imagine that a sends a message (of type μ) to b . (On this view, a can send at most one message to b). And if μ is invertible, then b replies by sending a message (of type μ^{-1}) back to a . If $\text{tp}^{\mathfrak{A}}[a, b]$ is silent, then neither element sends a message to the other.

The remainder of this section is devoted to three simple observations on structures interpreting counting signatures.

Definition 3. Let $\mathfrak{A}$ be a structure interpreting a counting signature Σ , and let Y be an integer. We say that $\mathfrak{A}$ is *Y -branching* (over Σ) if, for every $a \in A$ and every counting predicate f of Σ , the set $\{a' \in A \mid a \neq a' \text{ and } \mathfrak{A} \models f[a, a']\}$ has cardinality at most Y . We say that $\mathfrak{A}$ is *finitely branching* if $\mathfrak{A}$ is Y -branching for some $Y \in \mathbb{N}$.

Definition 4. Let $\mathfrak{A}$ be a structure interpreting a counting signature Σ . We say that $\mathfrak{A}$ is *chromatic* (over Σ) if, for all $a, a', a'' \in A$, the following two conditions hold:

1. if $a \neq a'$ and $\text{tp}^{\mathfrak{A}}[a, a']$ is an invertible message-type, then $\text{tp}^{\mathfrak{A}}[a] \neq \text{tp}^{\mathfrak{A}}[a']$;
2. if a, a', a'' are pairwise distinct and both $\text{tp}^{\mathfrak{A}}[a, a']$ and $\text{tp}^{\mathfrak{A}}[a', a'']$ are invertible message-types, then $\text{tp}^{\mathfrak{A}}[a] \neq \text{tp}^{\mathfrak{A}}[a'']$.

Thus, a chromatic structure is one in which distinct elements connected by a chain of 1 or 2 invertible message-types have distinct 1-types.

Lemma 2. *Let Σ be a counting signature with exactly m counting predicates, let Y be a non-negative integer, and let $\mathfrak{A}$ be a Y -branching structure interpreting Σ . Let Σ' be the signature formed by adding $\log((mY)^2 + 1)$ (rounded up) new unary predicates to Σ , and regard Σ' as a counting signature by taking the counting predicates of Σ' to be the same as the counting predicates of Σ . Then $\mathfrak{A}$ can be expanded to a chromatic structure $\mathfrak{A}'$ interpreting Σ' .*

Proof. Consider the (undirected) graph G on A whose edges are the pairs of distinct elements connected by a chain of 1 or 2 invertible message-types. That is, $G = (A, E^1 \cup E^2)$, where

$$E^1 = \{(a, a') \mid a \neq a' \text{ and } \text{tp}^{\mathfrak{A}}[a, a'] \text{ is an invertible message-type}\}$$

$$E^2 = \{(a, a'') \mid a \neq a'' \text{ and for some } a' \in A, (a, a') \text{ and } (a', a'') \text{ are both in } E^1\}.$$

Since there are only m counting predicates in Σ and $\mathfrak{A}$ is Y -branching, the degree of G (in the normal graph-theoretic sense) is at most $(mY)^2$. Now use the standard greedy algorithm to colour G with $(mY)^2 + 1$ colours. By interpreting the $\log((mY)^2 + 1)$ (rounded up) new unary predicates to encode these colours, we obtain the desired expansion $\mathfrak{A}'$. $\square$

Definition 5. Let $\mathfrak{A}$ be a structure and Z an integer. We say that $\mathfrak{A}$ is Z -differentiated if, for every 1-type π , the number u of elements in A having 1-type π satisfies either $u \leq 1$ or $u > Z$.

Lemma 3. *Let Σ be a counting signature, let $\mathfrak{A}$ be a structure interpreting Σ , and let Z be a positive integer. Let Σ' be the signature formed by adding $\log Z$ (rounded up) new unary predicates to Σ , and regard Σ' as a counting signature by taking the counting predicates of Σ' to be the same as the counting predicates of Σ . Then $\mathfrak{A}$ can be expanded to a Z -differentiated structure $\mathfrak{A}'$ interpreting Σ' . Moreover, if $\mathfrak{A}$ is chromatic over Σ , we can ensure that $\mathfrak{A}'$ is chromatic over Σ' .*

Proof. For each 1-type π realized more than once but no more than Z times, colour the elements having 1-type π using Z different colours. By interpreting the $\log Z$ (rounded up) new unary predicates to encode these colours, we obtain the desired expansion $\mathfrak{A}'$. This process clearly preserves chromaticity. $\square$

Definition 6. Let Σ be a counting signature, let $\mathfrak{A}$ be a structure interpreting Σ , and let π, π' be 1-types over Σ . We say that π and π' form a *noisy pair* in $\mathfrak{A}$ if there do not exist distinct $a, a' \in A$ such that $\text{tp}^{\mathfrak{A}}[a] = \pi$, $\text{tp}^{\mathfrak{A}}[a'] = \pi'$, and $\text{tp}^{\mathfrak{A}}[a, a']$ is silent.

Thus, if π and π' form a noisy pair, every element with 1-type π either sends a message to, or receives a message from, every element with 1-type π' . Note that Definition 6 does not require π and π' to be distinct.

Lemma 4. *Let Σ be a counting signature with exactly m counting predicates, let Y be a non-negative integer, let $\mathfrak{A}$ be a Y -branching structure interpreting Σ , and let π, π' be 1-types over Σ . If π and π' form a noisy pair in $\mathfrak{A}$, then either there are at most $(mY + 1)^2$ elements having 1-type π , or there are at most $(mY + 1)^2$ elements having 1-type π' .*

Proof. Suppose for contradiction that π and π' form a noisy pair, but that there are more than $(mY + 1)^2$ elements having 1-type π and more than $(mY + 1)^2$ elements having 1-type π' . Now let B be a set of elements having 1-type π , with $|B| > (mY)^2 + mY$; and let B' be a set of elements having 1-type π' , disjoint from B , with $|B'| = mY + 1$. Such sets can evidently be found even if $\pi = \pi'$. Select any $b \in B$. Since b sends a message to at most mY elements in B' , there exists $b' \in B'$ such that b sends no message to b' . Since π and π' form a noisy pair, we have that, for all $b \in B$, there exists $b' \in B'$ such that b' sends a message to b . But the total number of elements $b \in A$ such that b' sends a message to b for some $b' \in B'$ is bounded by $mY|B'| = mY(mY + 1)$, contradicting the supposition that $|B| > (mY)^2 + mY$. $\square$

3 Profiles and approximations

Let Σ be a finite counting signature with counting predicates $f_1, \dots, f_m$. Enumerate the invertible message-types over Σ (in some arbitrary order) as $\mu_1, \dots, \mu_{M^*}$, and the non-invertible message-types over Σ (again in some arbitrary order) as $\mu_{M^*+1}, \dots, \mu_M$. (Thus, $\mu_1, \dots, \mu_M$ is an enumeration of all the message-types over Σ .) Denote the set of silent 2-types over Σ by Ξ . We fix the symbols Σ , m , f_h ($1 \leq h \leq m$), M^* , M , μ_j ($1 \leq j \leq M$) and Ξ to have these meanings throughout this section.

We introduce the concepts of the Π -profile and the Π -count of an element in a structure interpreting Σ , where Π is a set of 1-types.

Notation 2. Let $\mathfrak{A}$ be a structure interpreting Σ , and let π be any 1-type and Π any set of 1-types over Σ . When $\mathfrak{A}$ is clear from context, denote by A_π the set $\{a \in A \mid \text{tp}^\mathfrak{A}[a] = \pi\}$, and denote by A_Π the set $\bigcup \{A_\pi \mid \pi \in \Pi\}$. In addition, denote by Π^c the set of all and only those 1-types over Σ not contained in Π .

Remark 2. For any $\mathfrak{A}$ interpreting Σ and any set of 1-types Π over Σ , $A_{\Pi^c} = A \setminus A_\Pi$.

For the next definition, recall that $\mu_1, \dots, \mu_M$ are the message-types (invertible and non-invertible) over Σ .

Definition 7. Let $\mathfrak{A}$ be a finitely branching structure interpreting Σ , let $a \in A$, and let Π be any set of 1-types over Σ . The Π -profile of a in $\mathfrak{A}$, denoted $\text{pr}_\Pi^\mathfrak{A}[a]$, is the M -element integer vector whose j th element ($1 \leq j \leq M$) is given by:

$$|\{b \in A_\Pi : b \neq a \text{ and } \text{tp}^\mathfrak{A}[a, b] = \mu_j\}|.$$

If Π is the set of all 1-types over Σ , we call $\text{pr}_\Pi^\mathfrak{A}[a]$ simply the *profile* of a in $\mathfrak{A}$, and denote it $\text{pr}^\mathfrak{A}[a]$.

Think of the vector $\text{pr}^{\mathfrak{A}}[a]$ as a description of the ‘local environment’ of a . The intuition is that a ’s local environment tells us, for each message-type μ , *how many* elements a is related to by μ ; but it tells us nothing about how those elements are related to each other, or about the elements of A to which a does not send a message. More generally, $\text{pr}_{\Pi}^{\mathfrak{A}}[a]$ is a partial description of that local environment—one that ignores elements whose 1-type is not in the set Π .

For the next definition, recall that $f_1, \dots, f_m$ are the counting predicates of Σ .

Definition 8. Let $\mathfrak{A}$ be a finitely branching structure interpreting Σ , let $a \in A$, and let Π be any set of 1-types over Σ . The Π -count of a in $\mathfrak{A}$, denoted $\text{ct}_{\Pi}^{\mathfrak{A}}[a]$, is the m -element integer vector whose h th element ($1 \leq h \leq m$) is given by:

$$|\{b \in A_{\Pi} : b \neq a \text{ and } \mathfrak{A} \models f_h[a, b]\}|.$$

Furthermore, a set $B \subseteq A$ is called a Π -group if every element of B has the same 1-type and every element of B has the same Π -count. If Π is the set of all 1-types over Σ , we call $\text{ct}_{\Pi}^{\mathfrak{A}}[a]$ simply the *count* of a in $\mathfrak{A}$, and denote it $\text{ct}^{\mathfrak{A}}[a]$.

Think of the vector $\text{ct}_{\Pi}^{\mathfrak{A}}[a]$ as providing a ‘statistical summary’ of the vector $\text{pr}_{\Pi}^{\mathfrak{A}}[a]$. In particular, $\text{pr}_{\Pi}^{\mathfrak{A}}[a]$ determines $\text{ct}_{\Pi}^{\mathfrak{A}}[a]$, but not conversely.

Definition 9. Let $\mathfrak{A}$ be a finitely branching, chromatic structure interpreting Σ , let Π be a set of 1-types over Σ , and let B be a subset of A . A structure $\mathfrak{A}'$ over the domain A is a (Π, B) -approximation to $\mathfrak{A}$ if (i) $\mathfrak{A}'$ is chromatic; (ii) every 2-type realized in $\mathfrak{A}'$ is also realized in $\mathfrak{A}$; and (iii) for all $a \in A$:

1. $\text{tp}^{\mathfrak{A}'}[a] = \text{tp}^{\mathfrak{A}}[a]$;
2. $\text{pr}_{\Pi^c}^{\mathfrak{A}'}[a] = \text{pr}_{\Pi^c}^{\mathfrak{A}}[a]$.
3. $a \in A \setminus B$ implies $\text{pr}^{\mathfrak{A}'}[a] = \text{pr}^{\mathfrak{A}}[a]$;
4. $a \in B$ implies $\text{ct}_{\Pi}^{\mathfrak{A}'}[a] = \text{ct}_{\Pi}^{\mathfrak{A}}[a]$.

In Condition 4 of the above definition, the restriction that $a \in B$ is in fact logically redundant; since if $a \notin B$, Condition 3 certainly entails $\text{ct}_{\Pi}^{\mathfrak{A}'}[a] = \text{ct}_{\Pi}^{\mathfrak{A}}[a]$.

Remark 3. If $\mathfrak{A}'$ is a (Π, B) -approximation to $\mathfrak{A}$, $\Pi \subseteq \Pi'$, and $B \subseteq B'$, then $\mathfrak{A}'$ is also a (Π', B') -approximation to $\mathfrak{A}$. If $\mathfrak{A}'$ is a (Π, B) -approximation to $\mathfrak{A}$, and $\mathfrak{A}''$ is a (Π, B) -approximation to $\mathfrak{A}'$, then $\mathfrak{A}''$ is a (Π, B) -approximation to $\mathfrak{A}$.

Our strategy now is to show that, in favourable circumstances, (Π, B) -approximations can be obtained in which the elements of B exhibit ‘few’ profiles. We first consider the special case where Π is a singleton. For the next definition, recall that $\mu_1, \dots, \mu_{M^*}$ are the invertible message-types over Σ ; thus, the first M^* coordinates of any π -profile $\text{pr}_{\{\pi\}}^{\mathfrak{A}}[a]$ tell us, for each invertible message-type μ , how many elements with 1-type π a sends a message of type μ to.

Definition 10. Let π be a 1-type over Σ , let $\mathfrak{A}$ be a finitely branching structure interpreting Σ , and let B be a subset of A . We say that B is a π -patch if B is a $\{\pi\}$ -group and, for all $a, b \in B$, the vectors $\text{pr}_{\{\pi\}}^{\mathfrak{A}}[a]$ and $\text{pr}_{\{\pi\}}^{\mathfrak{A}}[b]$ agree in each of their first M^* coordinates.

Lemma 5. *Let $\mathfrak{A}$ be a chromatic, Y -branching structure interpreting Σ , let π be a 1-type over Σ , and let $B \subseteq A$ be a π -patch. If $\mathfrak{A}$ is $(mY+1)^2$ -differentiated, then there exists a structure $\mathfrak{A}'$ such that $\mathfrak{A}'$ is a $(\{\pi\}, B)$ -approximation to $\mathfrak{A}$ in which the elements of B all have the same $\{\pi\}$ -profile.*

Proof. Let π^* be the 1-type such that $B \subseteq A_{\pi^*}$, and suppose that $\mathfrak{A}$ is $(mY+1)^2$ -differentiated. If $|B| \leq 1$, $\mathfrak{A}' = \mathfrak{A}$ obviously satisfies the conditions of the Lemma. And if $|A_{\pi}| \leq 1$, any two elements with the same $\{\pi\}$ -count have the same $\{\pi\}$ -profile, so that $\mathfrak{A}' = \mathfrak{A}$ again satisfies the conditions of the Lemma. So we may suppose that B and A_{π} both contain more than one element, whence $|A_{\pi^*}| > (mY+1)^2$ and $|A_{\pi}| > (mY+1)^2$. By Lemma 4, let τ be a silent 2-type such that, for some $a \in A_{\pi^*}$ and some $a' \in A_{\pi}$, $\text{tp}^{\mathfrak{A}}[a, a'] = \tau$.

For $a \in B$, let

$$A_a = \{a' \in A_{\pi} \mid a \neq a' \text{ and } \text{tp}^{\mathfrak{A}}[a, a'] \text{ is a non-invertible message-type}\};$$

and for $a \notin B$, let $A_a = \emptyset$. Notice, incidentally, that $a' \in A_a$ implies $a \notin A_{a'}$. Choose $b \in B$ for which $|A_b|$ is smallest, and fix b . Enumerate A_b (finite or infinite) as $b_1, b_2, \dots$. For any $a \in B$ not equal to b , let $\hat{A}_a$ be a subset of A_a having the same number of elements as A_b , and enumerate $\hat{A}_a$ as $a_1, a_2, \dots$.

We now define the structure $\mathfrak{A}'$ by assigning 2-types as follows. For all $a \in B$ such that $a \neq b$, set

$$\text{tp}^{\mathfrak{A}'}[a, a_i] = \text{tp}^{\mathfrak{A}}[b, b_i], \quad (1)$$

where i ranges over the enumeration of $\hat{A}_a$, and set

$$\text{tp}^{\mathfrak{A}'}[a, a'] = \tau, \quad (2)$$

where a' is any element of $A_a \setminus \hat{A}_a$. In addition, for all distinct a, a' such that $a' \notin A_a$ and $a \notin A_{a'}$, set

$$\text{tp}^{\mathfrak{A}'}[a, a'] = \text{tp}^{\mathfrak{A}}[a, a']. \quad (3)$$

Since $a' \in A_a$ implies $a \notin A_{a'}$, none of these assignments overwrites any other. And since $B \subseteq A_{\pi^*}$, the 1-type assignments implicit in (1)–(3) never clash: indeed, we have $\text{tp}^{\mathfrak{A}'}[a] = \text{tp}^{\mathfrak{A}}[a]$ for all $a \in A$. Furthermore, the transformation from $\mathfrak{A}$ to $\mathfrak{A}'$ does not affect invertible message-types. That is: for distinct a, a' , $\text{tp}^{\mathfrak{A}}[a, a']$ is an invertible message-type if and only if $\text{tp}^{\mathfrak{A}'}[a, a']$ is an invertible message-type; and moreover, if $\text{tp}^{\mathfrak{A}}[a, a']$ is an invertible message-type, then $\text{tp}^{\mathfrak{A}'}[a, a'] = \text{tp}^{\mathfrak{A}}[a, a']$.

We now verify that $\mathfrak{A}'$ is a $(\{\pi\}, B)$ -approximation to $\mathfrak{A}$. From the remarks of the previous paragraph and the fact that $\mathfrak{A}$ is chromatic, we have that $\mathfrak{A}'$

is also chromatic. In addition, it is immediate from (1)–(3) that every 2-type realized in $\mathfrak{A}'$ is also realized in $\mathfrak{A}$. Now let a be any element of B . Since B is a π -patch, the vectors $\text{pr}_{\{\pi\}}^{\mathfrak{A}}[a]$ and $\text{pr}_{\{\pi\}}^{\mathfrak{A}}[b]$ by definition agree in their first M^* coordinates (corresponding to the invertible message-types). Hence, since we have just shown that the transformation from $\mathfrak{A}$ to $\mathfrak{A}'$ does not affect invertible message-types, the vectors $\text{pr}_{\{\pi\}}^{\mathfrak{A}'}[a]$ and $\text{pr}_{\{\pi\}}^{\mathfrak{A}'}[b]$ also agree in their first M^* coordinates. Furthermore, the assignments (1) and (2) guarantee that $\text{pr}_{\{\pi\}}^{\mathfrak{A}'}[a]$ and $\text{pr}_{\{\pi\}}^{\mathfrak{A}'}[b]$ also agree in the remaining coordinates $M^* + 1, \dots, M$ (corresponding to the non-invertible message-types). Hence we have:

$$\text{pr}_{\{\pi\}}^{\mathfrak{A}'}[a] = \text{pr}_{\{\pi\}}^{\mathfrak{A}}[b] \quad \text{for all } a \in B. \quad (4)$$

It is now a simple matter to check the numbered conditions in Definition 9 (iii). Let $a \in A$.

1. We have already established that $\text{tp}^{\mathfrak{A}'}[a] = \text{tp}^{\mathfrak{A}}[a]$.
2. Let a' be any element of $A_{\{\pi\}^c}$ with $a \neq a'$. Then certainly $a' \notin A_a \subseteq A_\pi$, so that $\text{tp}^{\mathfrak{A}'}[a, a']$ can be different from $\text{tp}^{\mathfrak{A}}[a, a']$ only if $a \in A_{a'}$. But if $a \in A_{a'}$, then neither $\text{tp}^{\mathfrak{A}}[a, a']$ nor $\text{tp}^{\mathfrak{A}'}[a, a']$ can be a message-type. Hence, $\text{pr}_{\{\pi\}^c}^{\mathfrak{A}'}[a] = \text{pr}_{\{\pi\}^c}^{\mathfrak{A}}[a]$.
3. Suppose $a \in A \setminus B$, and let a' be any element of A with $a \neq a'$. The argument now proceeds much as for the previous condition: certainly, $a' \notin A_a = \emptyset$, so $\text{tp}^{\mathfrak{A}'}[a, a']$ can be different from $\text{tp}^{\mathfrak{A}}[a, a']$ only if $a \in A_{a'}$. But if $a \in A_{a'}$, then neither $\text{tp}^{\mathfrak{A}}[a, a']$ nor $\text{tp}^{\mathfrak{A}'}[a, a']$ can be a message-type. Hence $\text{pr}^{\mathfrak{A}'}[a] = \text{pr}^{\mathfrak{A}}[a]$.
4. If $a \in B$, Equation (4) yields $\text{ct}_{\{\pi\}}^{\mathfrak{A}'}[a] = \text{ct}_{\{\pi\}}^{\mathfrak{A}}[b]$. And since B is a $\{\pi\}$ -group, $\text{ct}_{\{\pi\}}^{\mathfrak{A}}[b] = \text{ct}_{\{\pi\}}^{\mathfrak{A}}[a]$. It follows that $\text{ct}_{\{\pi\}}^{\mathfrak{A}'}[a] = \text{ct}_{\{\pi\}}^{\mathfrak{A}}[a]$.

Finally, it is immediate from Equation (4) that all elements of B have the same $\{\pi\}$ -profile in $\mathfrak{A}'$. $\square$

Thus, Lemma 5 assures us that, as long as we are content to work with $(\{\pi\}, B)$ -approximations to highly differentiated structures, we may unify the $\{\pi\}$ -profiles of the elements in any π -patch B .

We next deal with the case where Π is not a singleton. Here, the possibilities for reducing the number of Π -profiles realized in a Π -group are more limited.

Lemma 6. *Let $\mathfrak{A}$ be a finitely branching structure interpreting Σ , let Π be a set of 1-types over Σ , let $B \subseteq A$ be a Π -group, and let ω be a permutation of B . Then there exists a structure $\mathfrak{A}'$ interpreting Σ over the domain A such that $\mathfrak{A}'$ is a (Π, B) -approximation to $\mathfrak{A}$, and for all $b \in B$, $\text{pr}_{\Pi}^{\mathfrak{A}'}[\omega(b)] = \text{pr}_{\Pi}^{\mathfrak{A}}[b]$.*

Proof. First, extend ω to the whole of A by setting $\omega(a) = a$ for $a \in A \setminus B$. Next, for all $b \in A$, define:

$$\omega_{b\Pi}(a) = \begin{cases} \omega(a) & \text{if } b \in A_\Pi \\ a & \text{otherwise;} \end{cases}$$

Thus, $\omega_{b\Pi}$ is a permutation of A (which may be the identity). Since $(\omega_{b\Pi})^{-1}$ and $(\omega^{-1})_{b\Pi}$ are the same permutation, we may unambiguously write $\omega_{b\Pi}^{-1}$. Clearly, ω fixes B setwise and $A \setminus B$ pointwise; so, therefore, does $\omega_{b\Pi}^{-1}$. Moreover, since B is a Π -group, either $B \subseteq A_\Pi$ or $B \subseteq A_{\Pi^c} = A \setminus A_\Pi$. Hence, ω fixes both A_Π and A_{Π^c} setwise, and so therefore does $\omega_{b\Pi}^{-1}$.

Define the structure $\mathfrak{A}'$ over domain A by setting, for all distinct $a, a' \in A$:

$$\text{tp}^{\mathfrak{A}'}[a, a'] = \text{tp}^{\mathfrak{A}}[\omega_{a'\Pi}^{-1}(a), \omega_{a\Pi}^{-1}(a')]. \quad (5)$$

To show that $\mathfrak{A}'$ is well-defined, we must show first that the elements $\omega_{a'\Pi}^{-1}(a)$ and $\omega_{a\Pi}^{-1}(a')$ in each instance of (5) are distinct, and second, that the 1-type assignments implicit in the different instances of (5) do not clash. Suppose, then that $a \neq a'$; we prove that $\omega_{a'\Pi}^{-1}(a) \neq \omega_{a\Pi}^{-1}(a')$. Since the permutations $\omega_{a\Pi}^{-1}$ and $\omega_{a'\Pi}^{-1}$ fix B setwise and $A \setminus B$ pointwise, we may assume that $a, a' \in B$. If $B \subseteq A_\Pi$, then $\omega_{a'\Pi}^{-1}(a) = \omega^{-1}(a)$ and $\omega_{a\Pi}^{-1}(a') = \omega^{-1}(a')$; if, on the other hand, $B \subseteq A \setminus A_\Pi$, then $\omega_{a'\Pi}^{-1}(a) = a$ and $\omega_{a\Pi}^{-1}(a') = a'$. Either way, $\omega_{a'\Pi}^{-1}(a) \neq \omega_{a\Pi}^{-1}(a')$. Next, we prove that the 1-type assignments in (5) never clash. Since all elements of B have the same 1-type, and since ω is the identity outside B , we have, for all a, a' , $\text{tp}^{\mathfrak{A}}[\omega_{a'\Pi}^{-1}(a)] = \text{tp}^{\mathfrak{A}}[a]$; thus, $\text{tp}^{\mathfrak{A}}[\omega_{a'\Pi}^{-1}(a)]$ does not depend on a' . Hence, the 1-type assignments implicit in (5) cannot clash, and $\mathfrak{A}'$ is indeed well-defined. In fact, this argument establishes that $\text{tp}^{\mathfrak{A}'}[a] = \text{tp}^{\mathfrak{A}}[a]$ for all $a \in A$.

We first check the numbered conditions of Definition 9 (iii) in turn. Let $a \in A$.

1. We have just established that $\text{tp}^{\mathfrak{A}'}[a] = \text{tp}^{\mathfrak{A}}[a]$.
2. For all $b \in A_{\Pi^c}$, $\omega_{b\Pi}^{-1}(a) = a$; in particular, if $a \in A_{\Pi^c}$, then $\omega_{a\Pi}^{-1}(a) = a$. Therefore, $\omega_{a\Pi}^{-1}$ is in fact a permutation of $A_{\Pi^c} \setminus \{a\}$, and moreover, for all $b \in A_{\Pi^c} \setminus \{a\}$, $\text{tp}^{\mathfrak{A}'}[a, b] = \text{tp}^{\mathfrak{A}}[a, \omega_{a\Pi}^{-1}(b)]$. Thus, the list of 2-types $\text{tp}^{\mathfrak{A}'}[a, b]$ obtained as b ranges over $A_{\Pi^c} \setminus \{a\}$ is (in some order) the list of 2-types $\text{tp}^{\mathfrak{A}}[a, b']$ obtained as b' ranges over $A_{\Pi^c} \setminus \{a\}$. It follows that $\text{pr}_{\Pi^c}^{\mathfrak{A}'}[a] = \text{pr}_{\Pi^c}^{\mathfrak{A}}[a]$.
3. If $a \in A \setminus B$, then, for all $b \in A$, $\omega_{b\Pi}^{-1}(a) = a$; in particular, $\omega_{a\Pi}^{-1}(a) = a$. Therefore, $\omega_{a\Pi}^{-1}$ is a permutation of $A \setminus \{a\}$, and moreover, for all $b \in A \setminus \{a\}$, $\text{tp}^{\mathfrak{A}'}[a, b] = \text{tp}^{\mathfrak{A}}[a, \omega_{a\Pi}^{-1}(b)]$. Thus, the list of 2-types $\text{tp}^{\mathfrak{A}'}[a, b]$ obtained as b ranges over $A \setminus \{a\}$ is (in some order) the list of 2-types $\text{tp}^{\mathfrak{A}}[a, b']$ obtained as b' ranges over $A \setminus \{a\}$. It follows that $\text{pr}^{\mathfrak{A}'}[a] = \text{pr}^{\mathfrak{A}}[a]$.
4. For all $b \in A_\Pi$, $\omega_{b\Pi}^{-1}(a) = \omega^{-1}(a)$; in particular, if $a \in A_\Pi$, then $\omega_{a\Pi}^{-1}(a) = \omega^{-1}(a)$. Therefore, $\omega_{a\Pi}^{-1}$ is a bijection from the set $A_\Pi \setminus \{a\}$ to the set $A_\Pi \setminus \{\omega^{-1}(a)\}$, and moreover, for all $b \in A_\Pi \setminus \{a\}$, $\text{tp}^{\mathfrak{A}'}[a, b] = \text{tp}^{\mathfrak{A}}[\omega^{-1}(a), \omega_{a\Pi}^{-1}(b)]$. Thus, the list of 2-types $\text{tp}^{\mathfrak{A}'}[a, b]$ obtained as b ranges over $A_\Pi \setminus \{a\}$ is (in some order) the list of 2-types $\text{tp}^{\mathfrak{A}}[\omega^{-1}(a), b']$ obtained as b' ranges over $A_\Pi \setminus \{\omega^{-1}(a)\}$. It follows that

$$\text{pr}_\Pi^{\mathfrak{A}'}[a] = \text{pr}_\Pi^{\mathfrak{A}}[\omega^{-1}(a)]. \quad (6)$$

Certainly, then, we have $\text{ct}_{\Pi}^{\mathfrak{A}'}[a] = \text{ct}_{\Pi}^{\mathfrak{A}}[\omega^{-1}(a)]$. But since B is a Π -group in $\mathfrak{A}$, $\text{ct}_{\Pi}^{\mathfrak{A}}[\omega^{-1}(a)] = \text{ct}_{\Pi}^{\mathfrak{A}}[a]$. Hence $\text{ct}_{\Pi}^{\mathfrak{A}'}[a] = \text{ct}_{\Pi}^{\mathfrak{A}}[a]$.

We have thus established that, for all $a \in A$, $\text{pr}_{\Pi}^{\mathfrak{A}'}[a] = \text{pr}_{\Pi}^{\mathfrak{A}}[\omega^{-1}(a)]$ and $\text{pr}_{\Pi^c}^{\mathfrak{A}'}[a] = \text{pr}_{\Pi^c}^{\mathfrak{A}}[a]$. Since $\mathfrak{A}$ is chromatic, it follows easily that $\mathfrak{A}'$ is chromatic and that all 2-types realized in $\mathfrak{A}'$ are realized in $\mathfrak{A}$. Hence, $\mathfrak{A}'$ is a (Π, B) -approximation to $\mathfrak{A}$. Finally, it follows by putting $a = \omega(b)$ in Equation (6) that, for all $b \in B$, $\text{pr}_{\Pi}^{\mathfrak{A}'}[\omega(b)] = \text{pr}_{\Pi}^{\mathfrak{A}}[b]$. $\square$

Let $\mathfrak{A}$, Π and B satisfy the conditions of Lemma 6. That lemma then assures us that, as long as we are content to work with (Π, B) -approximations, we may permute the Π -profiles of the elements in any Π -group B at will! To see the power of this idea, let $\mathfrak{A}$ be a structure, let Π' and Π'' be disjoint sets of 1-types, and let B be a subset of A whose elements realize J' different Π' -profiles and J'' different Π'' -profiles. If $\Pi = \Pi' \cup \Pi''$, how many Π -profiles are realized by the elements of B ? Answer: in general $J'J''$; however, the following lemma shows that, under certain circumstances, we can find a (Π, B) -approximation to $\mathfrak{A}$ for which the answer is smaller.

Lemma 7. *$\mathfrak{A}$ be a finitely branching structure interpreting Σ , let Π' , Π'' be disjoint, nonempty sets of 1-types, and let $\Pi = \Pi' \cup \Pi''$. Suppose $B \subseteq A$ is both a Π' -group and a Π'' -group, and hence also a Π -group. Let the number of different Π' -profiles realized in $\mathfrak{A}$ by the elements of B be J' ; and let the number of different Π'' -profiles realized in $\mathfrak{A}$ by the elements of B be J'' . Then there exists a (Π, B) -approximation $\mathfrak{A}''$ to $\mathfrak{A}$ in which at most $J' + J''$ different Π -profiles are realized by the elements of B .*

Proof. For perspicuity, we give the proof where B is finite; the modifications required for the general case are easy. Enumerate B as $b_1, \dots, b_I$. Let the various Π' -profiles realized by at least one element of B be $\bar{v}'_1, \dots, \bar{v}'_{J'}$; and let the various Π'' -profiles realized by at least one element of B be $\bar{v}''_1, \dots, \bar{v}''_{J''}$. Since B is a Π' -group, Lemma 6 guarantees that we can obtain a (Π', B) -approximation to $\mathfrak{A}$ in which the Π' -profiles of B are permuted at will. So let $\mathfrak{A}'$ be a (Π', B) -approximation to $\mathfrak{A}$ in which the Π' -profiles of the $b_1, \dots, b_I$ fall into consecutive blocks in the sense depicted in the middle column in Fig. 1. More precisely, we have integers $0 = I_1 < I_2 < \dots < I_{J'+1} = I$ such that, for all j ($1 \leq j \leq J'$), $\text{pr}_{\Pi'}^{\mathfrak{A}'}[b_i] = \bar{v}'_j$ for i in the range $[I_j + 1, I_{j+1}]$. Since B is also a Π'' -group, we can likewise obtain a structure $\mathfrak{A}''$ such that $\mathfrak{A}''$ is a (Π'', B) -approximation to $\mathfrak{A}'$ in which the elements of B have Π'' -profiles likewise arranged in consecutive blocks. Since $\mathfrak{A}''$ is a (Π'', B) -approximation to $\mathfrak{A}'$ and the sets Π' and Π'' are disjoint, the Π' -profiles of the elements of B will be unaffected by the transformation from $\mathfrak{A}'$ to $\mathfrak{A}''$: a typical alignment of Π' -profiles and Π'' -profiles in $\mathfrak{A}''$ is shown in Fig. 1. By inspection, at most $J' + J''$ Π -profiles are realized in $\mathfrak{A}''$ by the elements of B . From Remark 3, $\mathfrak{A}''$ is a (Π, B) -approximation to $\mathfrak{A}$, because $\mathfrak{A}'$ is a (Π', B) -approximation to $\mathfrak{A}$ and $\mathfrak{A}''$ is a (Π'', B) -approximation to $\mathfrak{A}'$. $\square$

This leads us to the main result of this section.

Element of B	Π' -profile in $\mathfrak{A}'$ (and also in $\mathfrak{A}''$)	Π'' -profile in $\mathfrak{A}''$
b_1	$\bar{v}'_1$	$\bar{v}''_1$
	$\bar{v}'_2$	$\bar{v}''_2$
	$\vdots$	$\vdots$
	$\bar{v}'_{J'}$	$\bar{v}''_{J''}$
b_I		

Figure 1: Arrangement of Π' -profiles and Π'' -profiles in B .

Lemma 8. *Let $\mathfrak{A}$ be a Y -branching, $(mY + 1)^2$ -differentiated, chromatic structure over Σ , let $l \geq 0$, let Π be a non-empty set of 1-types such that $|\Pi| \leq 2^l$, and let $B \subseteq A$ be a Π -group. Then there is a structure $\mathfrak{A}'$ such that $\mathfrak{A}'$ is a (Π, B) -approximation to $\mathfrak{A}$ and the number of different Π -profiles realized in $\mathfrak{A}'$ by the elements of B is at most $2^l(M^* + 1)(Y + 1)^{lm}$.*

Proof. By induction on l . To aid readability, let K_l stand for $2^l(M^* + 1)(Y + 1)^{lm}$. If $l = 0$, let $\Pi = \{\pi\}$. Decompose B into maximal π -patches $B_1, \dots, B_H$. Since $\mathfrak{A}$ is chromatic, the first M^* coordinates of $\text{pr}_{\{\pi\}}^{\mathfrak{A}}[a]$ are either all zero, or else are all zero except for a single occurrence of unity. Therefore, $H \leq M^* + 1$. Now let $\mathfrak{A}_0 = \mathfrak{A}$, and for all h ($1 \leq h \leq H$), apply Lemma 5 to obtain a structure $\mathfrak{A}_h$ such that $\mathfrak{A}_h$ is a $(\{\pi\}, B_h)$ -approximation to $\mathfrak{A}_{h-1}$ in which the elements of B_h all have the same $\{\pi\}$ -profile. By Remark 3, $\mathfrak{A}_H$ is a $(\{\pi\}, B)$ -approximation to $\mathfrak{A}$. And because the B_h are pairwise disjoint, $1 \leq h < h' \leq H$ implies $\text{pr}_{\{\pi\}}^{\mathfrak{A}_{h'}}[a] = \text{pr}_{\{\pi\}}^{\mathfrak{A}_h}[a]$ for all $a \in B_h$. Hence, the total number of $\{\pi\}$ -profiles realized by elements of B in $\mathfrak{A}_H$ is at most $H \leq M^* + 1 = K_0$. Thus, setting $\mathfrak{A}' = \mathfrak{A}_H$ establishes the case $l = 0$.

Now suppose $l > 0$. We may assume Π is not a singleton, since otherwise, we can employ the argument of the case $l = 0$; so let Π be partitioned into non-empty sets Π' and Π'' each of cardinality at most 2^{l-1} . Also, partition B into maximal Π' -groups $B_1, \dots, B_H$ (say). Since B is a Π -group and $\Pi' \cap \Pi'' = \emptyset$, the $B_1, \dots, B_H$ will also be Π'' -groups. Moreover, since $\mathfrak{A}$ is Y -branching, the Π -count of any element in $\mathfrak{A}$ is one of at most $(Y + 1)^m$ different vectors; and since B is a Π -group, all elements of B must have the same 1-type, whence $H \leq (Y + 1)^m$. Again, let $\mathfrak{A}_0 = \mathfrak{A}$, and consider the set B_1 . By inductive hypothesis, let $\mathfrak{A}'_1$ be a (Π', B_1) -approximation to $\mathfrak{A}_0$ in which at most K_{l-1} Π' -profiles are realized by the elements of B_1 . Again, by inductive hypothesis, let $\mathfrak{A}''_1$ be a (Π'', B_1) -approximation to $\mathfrak{A}'_1$ in which at most K_{l-1} Π'' -profiles are realized by the

elements of B_1 . Thus, in the structure $\mathfrak{A}'_1$, B_1 is a Π' -group realizing at most K_{l-1} different Π' -profiles, and also a Π'' -group realizing at most K_{l-1} different Π'' -profiles. By Lemma 7, let $\mathfrak{A}_1$ be a (Π, B_1) -approximation to $\mathfrak{A}'_1$ in which the elements of B_1 realize at most $2K_{l-1}$ different Π -profiles. By Remark 3, $\mathfrak{A}_1$ is a (Π, B_1) -approximation to $\mathfrak{A}_0$. Treating the sets $B_2, \dots, B_H$ in the same way, we obtain structures $\mathfrak{A}_h$ ($1 \leq h \leq H$) such that, for each h in this range, $\mathfrak{A}_h$ is a (Π, B) -approximation to $\mathfrak{A}_{h-1}$ in which at most $2K_{l-1}$ different Π -profiles are realized by the elements of B_h . By Remark 3, $\mathfrak{A}_H$ is a (Π, B) -approximation to $\mathfrak{A}$. And because the B_h are pairwise disjoint, $1 \leq h < h' \leq H$ implies that $\text{pr}^{\mathfrak{A}_{h'}}[a] = \text{pr}^{\mathfrak{A}_h}[a]$ for all $a \in B_h$. Hence, the total number of Π -profiles realized by elements of B in $\mathfrak{A}_H$ is at most

$$\begin{aligned} 2HK_{l-1} &\leq 2(Y+1)^m K_{l-1} \\ &= K_l. \end{aligned}$$

Thus, setting $\mathfrak{A}' = \mathfrak{A}_H$ completes the induction. $\square$

4 Deciding finite satisfiability

Let Σ be a signature of unary and binary predicates, let $\alpha(x)$ be a quantifier-free, equality-free formula over Σ with x as its only variable, let $\beta(x, y)$ be a quantifier-free, equality-free formula over Σ with x and y as its only variables, let m be a positive integer, let $f_1, \dots, f_m$ be binary predicates, let $C_1, \dots, C_m$ be positive integers, let ϕ^* be the formula

$$\phi^* := \forall x \alpha \wedge \forall x \forall y (\beta \vee x \approx y) \wedge \bigwedge_{1 \leq h \leq m} \forall x \exists =_{C_h} y (f_h(x, y) \wedge x \not\approx y), \quad (7)$$

and let $C = \max_{1 \leq h \leq m} C_h$. Make Σ into a counting signature by declaring the counting predicates of Σ to be exactly $\{f_1, \dots, f_m\}$. Let the number of symbols (unary and binary predicates) in Σ be s , so that the total number of 1-types over Σ is $L = 2^s$. As in Section 3, enumerate the invertible message-types over Σ (in some arbitrary order) as $\mu_1, \dots, \mu_{M^*}$, enumerate the non-invertible message-types over Σ (again in some arbitrary order) as $\mu_{M^*+1}, \dots, \mu_M$, and denote the set of silent 2-types over Σ by Ξ . We fix the symbols α, β, m, f_h ($1 \leq h \leq m$), C_h ($1 \leq h \leq m$), $C, \phi^*, \Sigma, s, L, M^*, M, \mu_j$ ($1 \leq j \leq M$) and Ξ to have these meanings throughout this section.

The motivation for introducing (Π, B) -approximations in Section 3 is that they are good enough for checking the (finite) satisfiability of ϕ^* .

Lemma 9. *Let $\mathfrak{A}$ be a finitely branching, chromatic structure interpreting Σ , let Π be a set of 1-types over Σ , let B be a subset of A , and let $\mathfrak{A}'$ be a (Π, B) -approximation to $\mathfrak{A}$. If $\mathfrak{A} \models \phi^*$, then $\mathfrak{A}' \models \phi^*$.*

Proof. By Remark 3, we may assume without loss of generality that Π is the set of all 1-types and $B = A$. Since $\text{tp}^{\mathfrak{A}'}[a] = \text{tp}^{\mathfrak{A}}[a]$ for every $a \in A$, $\mathfrak{A}' \models \forall x \alpha$. Since every 2-type realized in $\mathfrak{A}'$ is also realized in $\mathfrak{A}$, $\mathfrak{A}' \models \forall x \forall y (\beta \vee x \approx y)$. And

since $\text{ct}^{\mathfrak{A}'}[a] = \text{ct}^{\mathfrak{A}}[a]$ for every $a \in A$, $\mathfrak{A}' \models \bigwedge_{1 \leq h \leq m} \forall x \exists_{=C_h} y (f_h(x, y) \wedge x \not\approx y)$. $\square$

The next definition relies on conventions established in Notation 1.

Definition 11. A *star-type* (over Σ) is a pair $\sigma = \langle \pi, \bar{v} \rangle$, where π is a 1-type over Σ and $\bar{v} = (v_1, \dots, v_M)$ is a vector over $\mathbb{N}$ satisfying the condition that, for all j ($1 \leq j \leq M$), $v_j > 0$ implies $\text{tp}_1(\mu_j) = \pi$. We say that σ is *chromatic* if the set of integers

$$\{v_j \mid 1 \leq j \leq M^* \text{ and } \text{tp}_2(\mu_j) = \pi'\}$$

sums to either 0 or 1 for every 1-type π' , and sums to 0 in the case $\pi' = \pi$. If $\mathfrak{A}$ is a finitely branching structure interpreting Σ and $a \in A$, then $\langle \text{tp}^{\mathfrak{A}}[a], \text{pr}^{\mathfrak{A}}[a] \rangle$ is evidently a star-type, which we denote by $\text{st}^{\mathfrak{A}}[a]$. We say that the star-type σ is *realized* in $\mathfrak{A}$ if $\sigma = \text{st}^{\mathfrak{A}}[a]$ for some $a \in A$.

We note in passing that, if $\langle \pi, \bar{v} \rangle$ is a star-type, and $\bar{v}$ is not the zero-vector, then π is actually determined by $\bar{v}$.

Notation 3. If $\sigma = \langle \pi, \bar{v} \rangle$ is a star-type with $\bar{v} = (v_1, \dots, v_M)$, we denote π by $\text{tp}(\sigma)$ and v_j by $\sigma[j]$ for all j ($1 \leq j \leq M$).

Remark 4. If $\mathfrak{A}$ is a finitely branching structure interpreting Σ and $a \in A$, then $\text{tp}(\text{st}^{\mathfrak{A}}[a]) = \text{tp}^{\mathfrak{A}}[a]$. Moreover, $\mathfrak{A}$ is chromatic if and only if every star-type realized in $\mathfrak{A}$ is chromatic.

Remark 5. Let σ be a chromatic star-type and let j and j' be integers between 1 and M^* (so that μ_j and $\mu_{j'}$ are invertible message-types). If $\mu_j^{-1} = \mu_{j'}$, then either $\sigma[j] = 0$ or $\sigma[j'] = 0$. In particular, if $\mu_j^{-1} = \mu_j$, then $\sigma[j] = 0$.

Definition 12. Let $\mathfrak{A}$ be a finitely branching structure interpreting Σ , and let X be a positive integer. We say that $\mathfrak{A}$ is *X-sparse* if $\mathfrak{A}$ realizes no more than X different star-types—that is, if $|\{\text{st}^{\mathfrak{A}}[a] : a \in A\}| \leq X$.

For the next lemma, recall that s is the number of symbols in Σ , m is the number of counting predicates in Σ , and $C = \max_{1 \leq h \leq m} C_h$.

Lemma 10. Let $Z \geq (mC + 1)^2$, and let $X = 4^s(16^s + 1)(C + 1)^{sm}$. If there is a chromatic, Z -differentiated model of ϕ^* interpreting Σ , then there is a chromatic, Z -differentiated, X -sparse model of ϕ^* interpreting Σ over the same domain.

Proof. Suppose $\mathfrak{A}$ is a chromatic, Z -differentiated model of ϕ^* interpreting Σ . Since $\mathfrak{A} \models \phi^*$, $\mathfrak{A}$ is C -branching. Note that $M^* \leq 2^{4s}$. Let Π be the set of all 1-types over Σ ; thus, $|\Pi| = 2^s$. Let $A_1, \dots, A_H$ be a list of the nonempty sets A_π , where $\pi \in \Pi$; thus, $H \leq 2^s$. The sets A_h together partition A ; moreover, since $\mathfrak{A} \models \phi^*$, each A_h ($1 \leq h \leq H$) is a Π -group. Letting $\mathfrak{A}_0 = \mathfrak{A}$, by Lemma 8, we can obtain $\mathfrak{A}_1, \dots, \mathfrak{A}_H$ such that, for all h ($1 \leq h \leq H$), $\mathfrak{A}_h$ is a (Π, A_h) -approximation to $\mathfrak{A}_{h-1}$ in which at most $2^s(16^s + 1)(C + 1)^{sm}$ different profiles

are realized by the elements of A_h . Using by now familiar reasoning, $\mathfrak{A}_H$ is therefore a (Π, A) -approximation to $\mathfrak{A}$ realizing at most $H \cdot [2^s(16^s + 1)(C + 1)^{sm}] \leq 4^s(16^s + 1)(C + 1)^{sm} = X$ different star-types. By Lemma 9, $\mathfrak{A}_H \models \phi^*$. By Definition 9, $\mathfrak{A}_H$ is chromatic; and since $\text{tp}^{\mathfrak{A}_H}[a] = \text{tp}^{\mathfrak{A}}[a]$ for all $a \in A$, $\mathfrak{A}_H$ is also Z -differentiated. Thus $\mathfrak{A}' = \mathfrak{A}_H$ is the required structure. $\square$

Notation 4. We write $\mathcal{I}$ to denote the set of (unordered) pairs of (not necessarily distinct) integers between 1 and L . Formally: $\mathcal{I} = \{\{i, i'\} \mid 1 \leq i \leq i' \leq L\}$.

For the next definition, recall that Ξ is the set of silent 2-types over Σ .

Definition 13. A *frame* over Σ is a tuple $\mathcal{F} = (\bar{\sigma}, I, \theta)$, where $\bar{\sigma} = (\sigma_1, \dots, \sigma_N)$ is a list of pairwise distinct star-types over Σ , I is a subset of $\mathcal{I}$, and θ is a function $\theta : I \rightarrow \Xi$ such that, for all $\{i, i'\} \in I$ with $i \leq i'$, $\text{tp}_1(\theta(\{i, i'\})) = \pi_i$ and $\text{tp}_2(\theta(\{i, i'\})) = \pi_{i'}$. The *dimension* of $\bar{\sigma}$ is N . For Y a positive integer, $\mathcal{F}$ is *Y -bounded* if, for all k ($1 \leq k \leq N$) and all j ($1 \leq j \leq M$), $\sigma_k[j] \leq Y$. Finally, $\mathcal{F}$ is *chromatic* if every σ_k is chromatic.

Think of a frame $\mathcal{F}$ as a (putative) statistical summary of a structure $\mathfrak{A}$, specifically:

Definition 14. Let $\mathcal{F} = (\bar{\sigma}, I, \theta)$ be a frame over Σ , and let $\mathfrak{A}$ be a structure interpreting Σ . We say that $\mathcal{F}$ *describes* $\mathfrak{A}$ if the following conditions hold:

1. $\bar{\sigma}$ is a list of all and only those star-types realized in $\mathfrak{A}$;
2. I is the set of all and only those $\{i, i'\}$ ($1 \leq i \leq i' \leq L$) such that π_i and $\pi_{i'}$ do not form a noisy pair in $\mathfrak{A}$ (see Definition 6);
3. for each $\{i, i'\} \in I$, there exist $a \in A_\pi$ and $a' \in A_{\pi'}$ such that $a \neq a'$ and $\text{tp}^{\mathfrak{A}}[a, a'] = \theta(\{i, i'\})$.

Every finitely branching structure $\mathfrak{A}$ is evidently described by some (not necessarily unique) frame; and certain interesting properties of $\mathfrak{A}$ correspond to obvious properties of the frames which describe it, as we see from the following two lemmas.

Lemma 11. *Let $\mathfrak{A}$ be a finitely branching structure interpreting Σ and let $\mathcal{F}$ be a frame over Σ which describes $\mathfrak{A}$. Then: (i) $\mathfrak{A}$ is chromatic if and only if $\mathcal{F}$ is chromatic; (ii) $\mathfrak{A}$ is X -sparse if and only if $\mathcal{F}$ has dimension at most X ; and (iii) if $\mathfrak{A}$ is Y -branching then $\mathcal{F}$ has bound Y .*

Proof. Immediate. $\square$

For the next definition, recall that a 1-type π is simply a finite collection of formulas, so that $\bigwedge \pi$ denotes the conjunction of those formulas; similarly for 2-types.

Definition 15. Let $\mathcal{F} = (\bar{\sigma}, I, \theta)$ be a frame over Σ , where $\bar{\sigma} = (\sigma_1, \dots, \sigma_N)$. We write $\mathcal{F} \models \phi^*$ if the following conditions are satisfied:

1. for all k ($1 \leq k \leq N$) $\models \bigwedge \text{tp}(\sigma_k) \rightarrow \alpha$;
2. for all k ($1 \leq k \leq N$) and all j ($1 \leq j \leq M$), if $\sigma_k[j] > 0$ then $\models \bigwedge \mu_j \rightarrow \beta(x, y) \wedge \beta(y, x)$;
3. for all $\{i, i'\} \in I$, $\models \bigwedge \theta(\{i, i'\}) \rightarrow \beta(x, y) \wedge \beta(y, x)$;
4. for all k ($1 \leq k \leq N$) and all h ($1 \leq h \leq m$), the integers in the set $\{\sigma_k[j] \mid 1 \leq j \leq M \text{ and } f_h(x, y) \in \mu_j\}$ sum to C_h .

Remark 6. Let $\mathcal{F} = (\bar{\sigma}, I, \theta)$ be a frame over Σ , where $\bar{\sigma} = (\sigma_1, \dots, \sigma_N)$. If $\mathcal{F} \models \phi^*$, then, for all k ($1 \leq k \leq N$), $\sum_{1 \leq j \leq M} \sigma_k[j] \leq mC$.

Lemma 12. Let $\mathfrak{A}$ be a finitely branching structure interpreting Σ and let $\mathcal{F}$ be a frame over Σ describing $\mathfrak{A}$. If $\mathfrak{A} \models \phi^*$, then $\mathcal{F} \models \phi^*$.

Proof. Almost immediate. $\square$

However, while every finitely branching structure is described by some frame, not every frame describes a structure; and it is important for us to define a class of frames which do. Recall that L is the number of 1-types over Σ and M^* the number of invertible message-types over Σ .

Notation 5. Let $\mathcal{F} = (\bar{\sigma}, I, \theta)$ be a frame over Σ , where $\bar{\sigma} = (\sigma_1, \dots, \sigma_N)$. If $\mathcal{F}$ is clear from context, for integers i, j, k in the ranges $1 \leq i \leq L$, $1 \leq j \leq M^*$, $1 \leq k \leq N$ write:

$$\begin{aligned} o_{ik} &= \begin{cases} 1 & \text{if } \text{tp}(\sigma_k) = \pi_i \\ 0 & \text{otherwise;} \end{cases} \\ p_{ik} &= \begin{cases} 1 & \text{if, for all } j \text{ } (1 \leq j \leq M), \text{tp}_2(\mu_j) = \pi_i \text{ implies } \sigma_k[j] = 0 \\ 0 & \text{otherwise;} \end{cases} \\ q_{jk} &= \sigma_k[j]; \\ r_{ik} &= \sum \{\sigma_k[j] \mid M^* + 1 \leq j \leq M \text{ and } \text{tp}_2(\mu_j) = \pi_i\}; \\ s_{ik} &= \sum \{\sigma_k[j] \mid 1 \leq j \leq M \text{ and } \text{tp}_2(\mu_j) = \pi_i\}. \end{aligned}$$

Remark 7. Let $\mathcal{F}$ be a frame over Σ , let $\mathfrak{A}$ be a structure interpreting Σ , and suppose $\mathcal{F}$ describes $\mathfrak{A}$. In that case, the symbols o_{ik} , p_{ik} , q_{jk} , r_{ik} and s_{ik} in Notation 5 have the following interpretations with respect to $\mathfrak{A}$:

1. $o_{ik} = 1$ just in case every element with star-type σ_k has 1-type π_i ;
2. $p_{ik} = 1$ just in case no element with star-type σ_k sends a message to any element having 1-type π_i ;

3. q_{jk} counts how many messages of type μ_j any element having star-type σ_k sends;
4. r_{ik} is the total number elements of 1-type π_i to which any element having star-type σ_k sends a non-invertible message; and
5. s_{ik} is the total number elements of 1-type π_i to which any element having star-type σ_k sends a message.

With this notation in hand we can characterize a class of frames whose members are guaranteed to describe finite structures.

Definition 16. Let $\mathcal{F} = (\bar{\sigma}, I, \theta)$ be a frame over Σ , where $\bar{\sigma} = (\sigma_1, \dots, \sigma_N)$. Let $\bar{w} = (w_1, \dots, w_N)$ be a vector of positive integers. Using Notation 5, for all i ($1 \leq i \leq L$), all i' ($1 \leq i' \leq L$) and all j ($1 \leq j \leq M^*$), let:

$$u_i = \sum_{1 \leq k \leq N} o_{ik} w_k \quad v_j = \sum_{1 \leq k \leq N} q_{jk} w_k \quad x_{ii'} = \sum_{1 \leq k \leq N} o_{ik} p_{i'k} w_k.$$

Finally, let Z be a positive integer. We say that $\bar{w}$ is a Z -solution of $\mathcal{F}$ if the following conditions are satisfied for all i ($1 \leq i \leq L$), all i' ($1 \leq i' \leq L$), all j ($1 \leq j \leq M^*$) and all k ($1 \leq k \leq N$):

- (C1) $v_j = v_{j'}$, where j' is such that $\mu_j^{-1} = \mu_{j'}$;
- (C2) $s_{ik} \leq u_i$;
- (C3) $u_i \leq 1$ or $u_i > Z$;
- (C4) if $o_{ik} = 1$, then either $u_i > 1$ or $r_{i'k} \leq x_{i'i}$;
- (C5) if $\{i, i'\} \notin I$, then either $u_i \leq 1$ or $u_{i'} \leq 1$;
- (C6) if $\{i, i'\} \notin I$ and $o_{ik} = 1$, then either $u_i > 1$ or $r_{i'k} \geq x_{i'i}$.

Remark 8. Let $\mathcal{F}$ be as in Definition 16, and suppose that $\mathfrak{A}$ is a finite structure interpreting Σ such that $\mathfrak{A}$ is described by $\mathcal{F}$. For all k ($1 \leq k \leq N$), let w_k be the number of elements of A having star-type σ_k in $\mathfrak{A}$. In that case, the symbols u_i , v_j and $x_{ii'}$ in Definition 16 have the following interpretations with respect to $\mathfrak{A}$:

1. u_i is the number of elements of $a \in A$ such that $\text{tp}^{\mathfrak{A}}[a] = \pi_i$;
2. v_j is the number of pairs $\langle a, b \rangle \in A^2$ such that $a \neq b$ and $\text{tp}^{\mathfrak{A}}[a, b] = \mu_j$;
3. $x_{ii'}$ is the number of elements of $a \in A$ such that $\text{tp}^{\mathfrak{A}}[a] = \pi_i$ and a does not send a message to any element having 1-type $\pi_{i'}$.

The following Lemma shows that our definition of Z -correctness is not too stringent for the sorts of structures that interest us.

Lemma 13. *Let $\mathfrak{A}$ be a finite, Y -branching and Z -differentiated structure interpreting Σ , with $Z \geq (mY + 1)^2$. Let $\mathcal{F} = (\bar{\sigma}, I, \theta)$ be a frame over Σ . If $\mathcal{F}$ describes $\mathfrak{A}$, then $\mathcal{F}$ has a Z -solution.*

Proof. Let $\bar{\sigma} = (\sigma_1, \dots, \sigma_N)$, and let $w_k = |\{a \in A : \text{st}^{\mathfrak{A}}[a] = \sigma_k\}|$ for all k ($1 \leq k \leq N$). We show that $\bar{w} = (w_1, \dots, w_N)$ is a Z -solution of $\mathcal{F}$. In doing so, we make free use of Remarks 7 and 8. Note that, by construction, the $w_1, \dots, w_N$ are all positive.

C1: If $\mu_j^{-1} = \mu_{j'}$, then the sets $\{\langle a, b \rangle \mid a \neq b \text{ and } \text{tp}^{\mathfrak{A}}[a, b] = \mu_j\}$ and $\{\langle a, b \rangle \mid a \neq b \text{ and } \text{tp}^{\mathfrak{A}}[a, b] = \mu_{j'}\}$ can obviously be put in 1–1 correspondence, namely: $\langle a, b \rangle \mapsto \langle b, a \rangle$. But the cardinalities of these sets are v_j and $v_{j'}$, respectively.

C2: Since $\mathcal{F}$ describes $\mathfrak{A}$, any element of A having star-type σ_k sends a message to exactly s_{ik} elements having 1-type π_i . But u_i is the number of elements of A having 1-type π_i . Since σ_k is realized in $\mathfrak{A}$, $s_{ik} \leq u_i$.

C3: Immediate given that $\mathfrak{A}$ is Z -differentiated.

C4: If $o_{ik} = 1$ and $u_i \leq 1$, then $u_i = 1$, so that $\mathfrak{A}$ contains exactly one element with 1-type π_i ; moreover, this element has star-type σ_k . Denote this element by a . Thus, a sends a non-invertible message to exactly $r_{i'k}$ elements with 1-type $\pi_{i'}$. Clearly, none of these elements sends a message back to a (since otherwise a 's message to it would be invertible), so that there exist at least $r_{i'k}$ elements with 1-type $\pi_{i'}$ which do not send a message to a . But since a is the only element with 1-type π_i , there exist at least $r_{i'k}$ elements with 1-type $\pi_{i'}$ which do not send a message to any element of 1-type π_i . In other words, $r_{i'k} \leq x_{i'i}$.

C5: Since $\mathcal{F}$ describes $\mathfrak{A}$, $\{i, i'\} \notin I$ implies that the 1-types π_i and $\pi_{i'}$ form a noisy pair in $\mathfrak{A}$. In that case, by Lemma 4, either $u_i \leq (mY + 1)^2$ or $u_{i'} \leq (mY + 1)^2$. Since $\mathfrak{A}$ is Z -differentiated and $(mY + 1)^2 \leq Z$, $u_i \leq 1$ or $u_{i'} \leq 1$.

C6: As already observed, if $o_{ik} = 1$ and $u_i \leq 1$, then $\mathfrak{A}$ contains exactly one element a having 1-type π_i ; moreover, this element has star-type σ_k , and $x_{i'i}$ is the number of elements of 1-type $\pi_{i'}$ which do not send a message to a . Since $\mathcal{F}$ describes $\mathfrak{A}$, $\{i, i'\} \notin I$ implies that π_i and $\pi_{i'}$ form a noisy pair, whence a sends a message—in fact, a non-invertible message—to all $x_{i'i}$ of these elements. But since a has star-type σ_k , a sends a non-invertible message to exactly $r_{i'k}$ elements having 1-type $\pi_{i'}$. Thus, $r_{i'k} \geq x_{i'i}$. $\square$

We now prove a converse of Lemma 13.

Lemma 14. *Let $\mathcal{F}$ be a chromatic frame over Σ , and let $Z \geq 3mC - 1$. If $\mathcal{F}$ has a Z -solution and $\mathcal{F} \models \phi^*$, then there exists a finite structure $\mathfrak{A}$ interpreting Σ such that $\mathfrak{A} \models \phi^*$.*

Proof. Let $\mathcal{F} = (\bar{\sigma}, I, \theta)$, let $\bar{\sigma} = (\sigma_1, \dots, \sigma_N)$, and let $\bar{w} = (w_1, \dots, w_N)$ be a Z -solution of $\mathcal{F}$. In the sequel, we use the symbols $o_{ik}, p_{ik}, q_{jk}, r_{ik}$ and s_{ik} (with indices in the appropriate ranges), as specified in Notation 5, and the symbols

u_i , v_j and $x_{ii'}$ (again, with indices in the appropriate ranges), as specified in Definition 16. Hence, the conditions **C1–C6** of Definition 16 hold.

For every k ($1 \leq k \leq N$), let A_k be a set of cardinality w_k , and let A be the disjoint union of the A_k . Think of A_k as the set of elements which ‘want’ to have star-type σ_k . In addition, we define for all i ($1 \leq i \leq L$), all i' ($1 \leq i' \leq L$) and all j ($1 \leq j \leq M^*$):

$$\begin{aligned} U_i &= \bigcup \{A_k \mid 1 \leq k \leq N \text{ and } o_{ik} = 1\} \\ X_{ii'} &= \bigcup \{A_k \mid 1 \leq k \leq N \text{ and } o_{ik} p_{i'k} = 1\} \\ V_j &= \bigcup \{A_k \mid 1 \leq k \leq N \text{ and } q_{jk} = 1\}. \end{aligned}$$

Since $\mathcal{F}$ is chromatic, $q_{jk} \leq 1$ for all j ($1 \leq j \leq M^*$) and all k ($1 \leq k \leq N$). Thus, for all i , i' and j in the appropriate ranges:

$$|U_i| = u_i; \quad |X_{ii'}| = x_{ii'}; \quad |V_j| = v_j.$$

Think of U_i as the set of elements which ‘want’ to have 1-type π_i , $X_{ii'}$ as the set of elements in U_i which do not ‘want’ to send a message to any element in $U_{i'}$, and V_j as the set of elements which ‘want’ to send an (invertible) message of type μ_j to some other element. We remark that $A_k \subseteq U_i$ if and only if $\text{tp}(\sigma_k) = \pi_i$. Moreover, for all j ($1 \leq j \leq M^*$), there exists a unique i ($1 \leq i \leq L$) such that $V_j \subseteq U_i$ —namely, that i such that $\text{tp}_1(\mu_j) = \pi_i$. We now convert the domain A into a structure $\mathfrak{A}$ in four steps.

Step 1 (Interpreting the unary predicates and diagonals of binary predicates): For every k ($1 \leq k \leq N$) and every $a \in A_k$, set $\text{tp}^{\mathfrak{A}}[a] = \text{tp}(\sigma_k)$. At the end of this step, we have, for every i ($1 \leq i \leq L$) and every $a \in U_i$, $\text{tp}^{\mathfrak{A}}[a] = \pi_i$.

Step 2 (Fixing the invertible message-types): For every j ($1 \leq j \leq M^*$), let j' be such that $\mu_j^{-1} = \mu_{j'}$. By **C1**, V_j and $V_{j'}$ are equinumerous. If $j' > j$, pick some 1–1 correspondence between V_j and $V_{j'}$; and for every $a \in V_j$, set $\text{tp}^{\mathfrak{A}}[a, a'] = \mu_j$, where a' is the element of $V_{j'}$ corresponding to $a \in V_j$. This completes Step 2. We must show that these assignments are meaningful, do not clash with Step 1, and do not clash with each other. Suppose then that the assignment $\text{tp}^{\mathfrak{A}}[a, a'] = \mu_j$ is made, and that $\mu_j^{-1} = \mu_{j'}$. Thus, $a \in V_j$ and $a' \in V_{j'}$. To show that the assignment is meaningful, we must prove that $a \neq a'$. For contradiction, suppose $a = a'$, and let k be such that $a \in A_k$. But then $\sigma_k[j] > 0$ and $\sigma_k[j'] > 0$, which is impossible by Remark 5. To show that the assignment does not clash with Step 1, suppose $\mu_j^{-1} = \mu_{j'}$, and let i , i' be such that $V_j \subseteq U_i$ and $V_{j'} \subseteq U_{i'}$. As observed above, $\pi_i = \text{tp}_1(\mu_j)$ and $\pi_{i'} = \text{tp}_1(\mu_{j'}) = \text{tp}_2(\mu_j)$, which conforms to the assignments in Step 1. To show that these assignments do not clash with each other, it suffices to prove that, if $a \in V_j \cap V_h$, $a' \in V_{j'} \cap V_{h'}$, $\mu_j^{-1} = \mu_{j'}$ and $\mu_h^{-1} = \mu_{h'}$, then $j = h$. Suppose then that antecedent of this conditional holds; let k and k' be such that $a \in A_k$ and $a' \in A_{k'}$. Then $\sigma_{k'}[j'] > 0$ and $\sigma_{k'}[h'] > 0$. Since $\sigma_{k'}$ is a




Figure 2: Dealing with non-invertible messages.

star-type, $\text{tp}_1(\mu_{j'}) = \text{tp}_1(\mu_{h'})$, whence $\text{tp}_2(\mu_j) = \text{tp}_2(\mu_h)$. But $\sigma_k[j] > 0$ and $\sigma_k[h] > 0$, and since σ_k is a *chromatic* star-type, $j = h$. Note that, if $\mu_j^{-1} = \mu_j$, then $V_j = \emptyset$ by Remark 5. Thus, at the end of Step 2, for every element $a \in A$ and every j ($1 \leq j \leq M^*$), a sends a (unique) message of type μ_j to some other element if and only if $a \in V_j$. That is: for all k ($1 \leq k \leq N$), all $a \in A_k$, and all j ($1 \leq j \leq M^*$), there are exactly $\sigma_k[j]$ elements $a' \in A$ such that $a \neq a'$ and $\text{tp}^{\mathfrak{A}}[a, a']$ has been set to the (invertible) message-type μ_j . We make one further observation before proceeding. Suppose that $\text{tp}^{\mathfrak{A}}[a, a']$ is assigned in this step and that $a \in U_i$; we claim that $a' \notin X_{i'i}$ for any i' . To see this, suppose $a \in A_k \subseteq V_j$ and $a' \in A_{k'} \subseteq V_{j'}$, with $\mu_j^{-1} = \mu_{j'}$. If $a \in U_i$, then $\text{tp}_1(\mu_j) = \text{tp}_2(\mu_{j'}) = \pi_i$. But $\sigma_{k'}[j] > 0$, whence $p_{ik'} = 0$, whence $a' \notin X_{i'i}$. This observation will be useful in Step 3.

Step 3 (Fixing the non-invertible message-types): Let i and i' be such that $1 \leq i \leq i' \leq L$. We fix all the non-invertible messages sent, in either direction, between U_i and $U_{i'}$. By **C3**, either $u_i \leq 1$ or $u_i > Z$; similarly, either $u_{i'} \leq 1$ or $u_{i'} > Z$. We consider five cases.

Case 1: $u_i = 0$. In this case, there are no elements of U_i and hence no 2-type assignments to be made between elements of U_i and elements of $U_{i'}$. Note that, by **C2**, $s_{ik} = 0$ for all k ($1 \leq k \leq N$), whence $\sigma_k[j] = 0$ for all k ($1 \leq k \leq N$) and for all j ($1 \leq j \leq M$) such that $\text{tp}_2(\mu_j) = \pi_i$. (Intuitively, no element of A —and in particular of $U_{i'}$ —‘wants’ to send a message to an element with 1-type π_i anyway.)

Case 2: $u_i = 1$. Let a be the sole element of U_i , and let k be such that $a \in A_k$. We deal first with the assignment of non-invertible messages sent from $U_{i'}$ to $U_i = \{a\}$. Consider any $a' \in A_{k'} \subseteq U_{i'}$. By **C2**, $s_{ik'} \leq 1$; hence there is at most one value of j in the range $1 \leq j \leq M$ such that $\text{tp}_2(\mu_j) = \pi_i$ and $\sigma_{k'}[j] > 0$. Suppose then that such a j exists. Again, since $s_{ik'} \leq 1$,

$\sigma_{k'}[j] = 1$. If $j \leq M^*$, then this message has already been dealt with in Step 2; otherwise, set $\text{tp}^{\mathfrak{A}}[a', a] = \mu_j$. Since $\text{tp}_1(\mu_j) = \pi_{i'}$ and $\text{tp}_2(\mu_j) = \pi_i$, this assignment does not clash with Step 1. Observe also that, just as in Step 2, if this assignment is made, we have, by definition, $p_{ik'} = 0$, so that $a' \notin X_{i'i}$. By carrying out the same procedure for all $a' \in U_{i'}$, we complete the assignment of non-invertible messages sent from $U_{i'}$ to U_i . It remains to deal with the non-invertible messages sent from $U_i = \{a\}$ to $U_{i'}$. Remembering that $a \in A_k$, **C4** ensures the existence of a subset $R \subseteq X_{i'i}$ such that $|R| = r_{i'k}$. For each j ($M^* + 1 \leq j \leq M$), if $\text{tp}_2(\mu_j) = \pi_{i'}$, select q_{jk} fresh elements a' of R , and make the assignment $\text{tp}^{\mathfrak{A}}[a, a'] = \mu_j$. (There are enough such elements by the definition of $r_{i'k}$.) Since $\text{tp}_1(\mu_j) = \pi_i$ and $\text{tp}_2(\mu_j) = \pi_{i'}$, this assignment does not clash with Step 1. Moreover, we have observed that $\text{tp}^{\mathfrak{A}}[a, a']$ has previously been assigned (either in this step or in Step 2) only if $a' \notin X_{i'i}$. Thus, these assignments do not clash with those made earlier in this step or those made in Step 2. The situation is illustrated in Fig. 2 Case 2.

Case 3: $u_{i'} = 0$ and $u_i > Z$. Symmetrical to Case 1.

Case 4: $u_{i'} = 1$ and $u_i > Z$. Symmetrical to Case 2.

Case 5: $u_i > Z$ and $u_{i'} > Z$. Since $Z \geq 3mC - 1$, partition U_i into three sets U_{i0}, U_{i1}, U_{i2} , each containing at least mC elements; and similarly for $U_{i'}$. Suppose $a \in U_i$. Then for some h ($0 \leq h < 3$), $a \in U_{ih}$. Let k be such that $a \in A_k$, and let $h' = h + 1 \pmod{3}$. For all j ($M^* + 1 \leq j \leq M$), pick $\sigma_k[j]$ fresh elements a' of $U_{i'h'}$ such that $\text{tp}^{\mathfrak{A}}[a, a']$ was not assigned in Step 2, and set $\text{tp}^{\mathfrak{A}}[a, a'] = \mu_j$. By Remark 6, we have, for each k ($1 \leq k \leq N$), $\sum_{1 \leq j \leq M} \sigma_k[j] \leq mC$; and since $|U_{i'h'}| \geq mC$, we never run out of fresh elements to pick. In this way, we deal with all messages sent from U_i to $U_{i'}$; the messages sent from $U_{i'}$ to U_i are dealt with symmetrically. It is obvious that these assignments do not clash with Step 1 (or Step 2); and the fact that $h' = h + 1 \pmod{3}$, ensures that they do not clash with each other (even if $i = i'$), as is evident from Fig. 2 Case 5.

Performing these assignments for all pairs i, i' such that $1 \leq i \leq i' \leq L$ completes Step 3. At the end of Step 3, then, for all k ($1 \leq k \leq N$), all $a \in A_k$, and all j ($1 \leq j \leq M$), there are exactly $\sigma_k[j]$ elements $a' \in A$ such that $a \neq a'$ and $\text{tp}^{\mathfrak{A}}[a, a'] = \mu_j$.

Step 4 (Fixing the silent 2-types): Finally, we deal with pairs of distinct elements a, a' whose 2-type in $\mathfrak{A}$ has not been yet assigned. Let i, i' be such that $a \in U_i$ and $a' \in U_{i'}$, and suppose, without loss of generality, that $i \leq i'$. We claim that $\langle i, i' \rangle \in I$. For suppose otherwise. By **C5**, we have either $u_i = 1$ or $u_{i'} = 1$. Assume the former. Now let k, k' be such that $a \in A_k$ and $a' \in A_{k'}$. Thus, $o_{i'k'} = 1$. If $p_{ik'} = 0$ then there is some j ($1 \leq j \leq M$) such that $q_{jk'} = \sigma_{k'}[j] > 0$ and $\text{tp}_2(\mu_j) = \pi_i$, whence—bearing in mind that a is the unique element of U_i — $\text{tp}^{\mathfrak{A}}[a, a']$ will certainly have been assigned in Step 2 (if μ_j is an invertible message-type) or in Step 3 Case 2 (if μ_j is a non-invertible message-type), contradicting our supposition. So we may assume that

$p_{ik'} = 1$ and hence $o_{i'k'}p_{ik'} = 1$. That is: $a \in X_{i'i}$. But $|X_{i'i}| = x_{i'i}$. And by **C6**, $x_{i'i} \leq r_{i'k}$. Yet in Step 3 (Case 2), $r_{i'k}$ elements of $X_{i'i}$ were chosen to receive messages from a . Hence a' must be among these elements, so that $\text{tp}^{\mathfrak{A}}[a, a']$ has already been assigned, again a contradiction. The case where $u_{i'} \leq 1$ proceeds symmetrically. Thus, we have established that, if $\text{tp}^{\mathfrak{A}}[a, a']$ has not yet been assigned, then $\{i, i'\} \in I$, so that we can make the assignment $\text{tp}^{\mathfrak{A}}[a, a'] = \theta(\{i, i'\})$. Since $\text{tp}_1(\theta(\{i, i'\})) = \pi_i$ and $\text{tp}_2(\theta(\{i, i'\})) = \pi_{i'}$, there is no clash with Step 1. Evidently, we can proceed in this way until all remaining 2-types have been assigned.

This completes the construction of $\mathfrak{A}$. The only 1-types realized in $\mathfrak{A}$ are the 1-types $\text{tp}(\sigma_k)$ (where $1 \leq k \leq N$). The only message-types realized in $\mathfrak{A}$ are those μ_j such that $\sigma_k[j] > 0$ for some k . And the only silent 2-types realized in $\mathfrak{A}$ are the $\theta(\{i, i'\})$ for $\{i, i'\} \in I$. Since $\mathcal{F} \models \phi^*$, we have $\mathfrak{A} \models \forall x \alpha \wedge \forall x \forall y (\beta \vee x \approx y)$. Moreover, for all k ($1 \leq k \leq N$), and for all $a \in A_k$, we have $\text{st}^{\mathfrak{A}}[a] = \sigma_k$. Since $\mathcal{F} \models \phi^*$, we have, for all $a \in A$, $\text{ct}^{\mathfrak{A}}[a] = (C_1, \dots, C_m)$; in other words, $\mathfrak{A} \models \bigwedge_{1 \leq h \leq m} \forall x \exists_{=C_h} y (f_h(x, y) \wedge x \not\approx y)$. Hence, $\mathfrak{A} \models \phi^*$. $\square$

We can now employ a standard result to bound the complexity of determining whether a given frame has a Z -solution.

Lemma 15. *Let $\mathcal{F}$ be a Y -bounded, N -dimensional frame over Σ , and let Z be an integer. Then $\mathcal{F}$ has a Z -solution if and only if it has a Z -solution $\bar{w}$ such that every component of $\bar{w}$ is bounded by some (fixed) singly exponential function of the quantity $L + M^* + N + \log Y + \log Z$.*

Proof. This follows immediately from the well-known result (Papadimitriou [6]) that, if an integer programming problem has a solution at all, then it has a solution all of whose components are bounded by a singly exponential function of the size of the problem (encoded in the obvious way). But the conditions **C1–C6** in Definition 16 amount to a disjunction of integer programming problems whose sizes are all bounded by a polynomial function of $L + M^* + N + \log Y + \log Z$. $\square$

Theorem 1. *The problem $\text{Fin-Sat-}\mathcal{C}^2$ is in NEXPTIME .*

Proof. Given ϕ , let $m, f_1, \dots, f_m, C_1, \dots, C_m$ and ϕ^* be as in Lemma 1, and let $C = \max_{1 \leq h \leq m} C_h$; thus $C \leq 2^{\|\phi^*\|}$. Let $Z = \max(3mC - 1, (mC + 1)^2)$. Let Σ be the signature of ϕ^* together with $\log((mC)^2 + 1) + \log Z$ (rounded up) additional unary predicates, regarded as a counting signature by taking the counting predicates of Σ to be $f_1, \dots, f_m$. Denote the total size of Σ by s : evidently, s is bounded by a polynomial function of $\|\phi^*\|$. Finally, let $X = 4^s \cdot (16^s + 1)(C + 1)^{sm}$.

Since $C \leq 2^{\|\phi^*\|}$, C is bounded by a singly exponential function of $\|\phi\|$. Thus, the problem of determining the satisfiability of ϕ over structures of size C or less is obviously in NEXPTIME . By Lemma 1 then, it suffices to show that the satisfiability of ϕ^* can be decided nondeterministically in time bounded by a singly exponential function of $\|\phi^*\|$.

We claim that ϕ^* is finitely satisfiable if and only if there exists a chromatic, C -bounded frame $\mathcal{F}$ over Σ of dimension $N \leq X$, such that $\mathcal{F}$ has a Z -solution and $\mathcal{F} \models \phi^*$. For suppose ϕ^* is finitely satisfiable. By Lemmas 2 and 3, ϕ^* has a finite, chromatic, Z -differentiated model $\mathfrak{A}$ interpreting Σ . By Lemma 10, we may assume without loss of generality that $\mathfrak{A}$ is also X -sparse. Let $\mathcal{F}$ be a frame over Σ describing $\mathfrak{A}$. By Lemma 11, $\mathcal{F}$ is chromatic, of dimension $N \leq X$ and C -bounded. By Lemma 12, $\mathcal{F} \models \phi^*$. Since $Z \geq (mC + 1)^2$, Lemma 13 guarantees that $\mathcal{F}$ has a Z -solution. Conversely, suppose $\mathcal{F}$ is a chromatic frame over Σ such that $\mathcal{F}$ has a Z -solution and $\mathcal{F} \models \phi^*$. Since $Z \geq 3mC - 1$, Lemma 14 guarantees that ϕ^* is finitely satisfiable.

By Lemma 15, then, ϕ^* is satisfiable if and only if there exists a chromatic, C -bounded frame $\mathcal{F}$ over Σ of dimension $N \leq X$ and a vector $\bar{w}$ of positive integers bounded by some doubly exponential function of $\|\phi^*\|$, such that $\mathcal{F} \models \phi^*$ and $\bar{w}$ is a Z -solution of $\mathcal{F}$. Using the standard binary encoding of integers, it is easy to write down $\mathcal{F}$ and $\bar{w}$ in a number of bits bounded by a singly exponential function of $\|\phi^*\|$, and to check whether they satisfy the above conditions in time bounded by a singly exponential function of $\|\phi^*\|$. This completes the proof. $\square$

We note in passing that the above proof yields a small model property for finitely satisfiable $\mathcal{C}^2$ -formulas.

Corollary 1. *Let ϕ be a formula of $\mathcal{C}^2$. If ϕ is finitely satisfiable, then it is satisfiable in a structure of size bounded by a doubly exponential function of $\|\phi\|$.*

Proof. The structure built in Lemma 14 from $\mathcal{F}$ and its Z -solution $\bar{w}$ has domain of cardinality $w_1 + \dots + w_N$. $\square$

Notice that the complexity result of Theorem 1 is better than one might naïvely expect on the basis of the small model property of Corollary 1. Nevertheless, the bound of Corollary 1 is optimal in the sense that there is a sequence $\{\phi_i\}$ of finitely satisfiable formulas of $\mathcal{C}^2$ whose size grows as a polynomial function of i , but whose smallest satisfying structures grow as a doubly exponential function of i (Grädel and Otto [2], p. 317).

5 Deciding satisfiability

Having established the complexity of determining finite satisfiability in $\mathcal{C}^2$, we now turn to the corresponding (general) satisfiability problem. In fact, there is almost no further work to do.

Notation 6. Let $\mathbb{N}^*$ denote the set $\mathbb{N} \cup \{\aleph_0\}$. We extend the ordering $>$ and the arithmetic operations $+$ and $\cdot$ from $\mathbb{N}$ to $\mathbb{N}^*$ in the obvious way. Specifically, we define $\aleph_0 > n$ for all $n \in \mathbb{N}$; we define $\aleph_0 + \aleph_0 = \aleph_0 \cdot \aleph_0 = \aleph_0$ and $0 \cdot \aleph_0 = \aleph_0 \cdot 0 = 0$; we define $n + \aleph_0 = \aleph_0 + n = \aleph_0$ for all $n \in \mathbb{N}$; and we define $n \cdot \aleph_0 = \aleph_0 \cdot n = \aleph_0$ for all $n \in \mathbb{N}$ such that $n > 0$. Under this extension, $>$ remains a total order, and $+$, $\cdot$ remain associative and commutative.

A system of linear equalities and inequalities defining an integer programming problem can of course be re-interpreted so that solutions are sought not over $\mathbb{N}$ but over $\mathbb{N}^*$. (We always assume that the coefficients occurring in such problems are in $\mathbb{N}$.) As an example, the single inequality $x_1 \geq x_1 + 1$ has no solutions over $\mathbb{N}$, but it does have a solution over $\mathbb{N}^*$, namely, $x_1 = \aleph_0$.

Lemmas 2–10 apply to both finite and infinite structures. Furthermore, the definition of a frame and its relationship to the (finitely branching) structures it describes make no reference to the cardinalities of those structures, and Lemmas 11–12 again apply generally. The concept of a Z -solution introduced in Definition 16 requires extension, however.

Definition 17. Let Σ and $\mathcal{F} = (\bar{\sigma}, I, \theta)$ be as in Definition 16, and let Z be a positive integer. Let $\bar{w} = (w_1, \dots, w_N)$ be a vector of non-zero elements of $\mathbb{N}^*$. We say that $\bar{w}$ is an *extended Z -solution* of $\mathcal{F}$ if $\bar{w}$ satisfies the conditions of Definition 16, with the arithmetic interpreted over $\mathbb{N}^*$ as specified in Notation 6.

We must check that the obvious analogues of Lemmas 13 and 14 hold:

Lemma 16. *Let $\mathfrak{A}$ be a Y -branching and Z -differentiated structure interpreting Σ , with $Z \geq (mY + 1)^2$. Let $\mathcal{F} = (\bar{\sigma}, I, \theta)$ be a frame over Σ . If $\mathcal{F}$ describes $\mathfrak{A}$, then $\mathcal{F}$ has an extended Z -solution.*

Lemma 17. *Let $\mathcal{F}$ be a chromatic frame over Σ , and let $Z \geq 3mC - 1$. If $\mathcal{F}$ has an extended Z -solution and $\mathcal{F} \models \phi^*$, then there exists a structure $\mathfrak{A}$ interpreting Σ such that $\mathfrak{A} \models \phi^*$.*

The proofs are exactly the same as in the finite case. Note that the variables u_i , v_j and $x_{ii'}$ as well as the w_k may now take the value $\aleph_0$; by contrast, the coefficients o_{ik} , p_{ik} , q_{jk} , r_{ik} and s_{ik} remain finite. Remark 8 generalizes unproblematically to countably infinite structures, so that the quantities u_i , v_j and $x_{ii'}$ mentioned in Definition 16 continue to have their familiar interpretations. The proofs of Lemmas 16 and 17 then proceed exactly as before.

There is one final hurdle to overcome. The proof of Lemma 15 used a well-known result bounding solutions of integer programming problems. Since we are dealing with $\mathbb{N}^*$ -programming problems, we need the following extension of that result.

Lemma 18. *Let Φ be a finite set of linear inequalities of the form*

$$a_0 + a_1x_1 + \dots + a_nx_n \leq b_0 + b_1x_1 + \dots + b_nx_n$$

in variables $x_1, \dots, x_n$. Here, all coefficients are assumed to be in $\mathbb{N}$. We take the size of Φ , denoted $\|\Phi\|$, to be measured in the usual way, assuming binary encoding of integers. If Φ has a solution over $\mathbb{N}^$, then Φ has a solution over $\mathbb{N}^*$ such that all finite values are bounded by some (fixed) singly exponential function of $\|\Phi\|$.*

Proof. Suppose that Φ has a solution over $\mathbb{N}^*$. Re-order the variables if necessary so that this solution has the form $\bar{a}\bar{\aleph}_0$, with $\bar{a} = a_1, \dots, a_k \in \mathbb{N}^k$ for some k ($0 \leq k \leq n$) and $\bar{\aleph}_0$ a $(n - k)$ -tuple of $\aleph_0$ s. Say that an inequality in Φ *does not involve* the variable x_i if the corresponding coefficients a_i and b_i are both zero. Let Ψ be the set of inequalities in Φ involving none of the $x_{k+1}, \dots, x_n$. Thus, Ψ , considered as a problem in variables $x_1, \dots, x_k$, has a solution $\bar{a}$ over $\mathbb{N}$, whence it has a solution $\bar{a}'$ bounded by some singly exponential function of $\|\Psi\|$ (and hence of $\|\Phi\|$). But then it is easy to see that $\bar{a}'\bar{\aleph}_0$ is a solution of Φ . $\square$

Theorem 2. *The problem $\text{Sat-}\mathcal{C}^2$ is in NEXPTIME.*

Proof. Exactly as for Theorem 1, noting that, by Lemma 18, the existence of extended Z-solutions for a frame $\mathcal{F}$ can be checked nondeterministically in time bounded by an exponential function of $\|\phi^*\|$. $\square$

Obviously, there is no interesting small model property for satisfiable $\mathcal{C}^2$ -formulas corresponding to Corollary 1; however, we have the next best thing.

Corollary 2. *Let ϕ be a formula of $\mathcal{C}^2$. Then there exist integers X and W , with X bounded by a singly exponential function of $\|\phi\|$ and W by a doubly exponential function of $\|\phi\|$ such that, if ϕ is satisfiable, then it is satisfiable in an X -sparse structure in which every star-type is realized either infinitely often or at most W times.*

References

- [1] Erich Grädel and Martin Otto. On logics with two variables. *Theoretical Computer Science*, 224(1–2):73–113, 1999.
- [2] Erich Grädel, Martin Otto, and Eric Rosen. Two-variable logic with counting is decidable. In *Proceedings of the 12th IEEE Symposium on Logic in Computer Science*, pages 306–317. IEEE Online Publications, 1997.
- [3] Martin Otto. Two-variable first-order logic over ordered domains. *Journal of Symbolic Logic*, 66(2):685–702, 2001.
- [4] Leszek Pacholski, Wiesław Szwaast, and Lidia Tendera. Complexity of two-variable logic with counting. In *Proceedings of the 12th IEEE Symposium on Logic in Computer Science*, pages 318–327. IEEE Online Publications, 1997.
- [5] Leszek Pacholski, Wiesław Szwaast, and Lidia Tendera. Complexity results for first-order two-variable logic with counting. *SIAM Journal on Computing*, 29(4):1083–1117, 1999.
- [6] Christos H. Papadimitriou. On the complexity of integer programming. *Journal of the Association for Computing Machinery*, 28(4):765–768, 1981.